

## Department of Veterans Affairs

## § 3.375

qualify as active tuberculosis. The requirements for presumptive service connection will be the same as those for tuberculous pleurisy.

(c) *Tuberculous pleurisy and endobronchial tuberculosis.* Tuberculous pleurisy and endobronchial tuberculosis fall within the category of pulmonary tuberculosis for the purpose of service connection on a presumptive basis. Either will be held incurred in service when initially manifested within 36 months after the veteran's separation from service as determined under § 3.307(a)(2).

(d) *Miliary tuberculosis.* Service connection for miliary tuberculosis involving the lungs is to be determined in the same manner as for other active pulmonary tuberculosis.

[26 FR 1591, Feb. 24, 1961, as amended at 27 FR 6387, July 6, 1962; 31 FR 4681, Mar. 19, 1966; 39 FR 34532, Sept. 26, 1974; 43 FR 45347, Oct. 2, 1978; 54 FR 34981, Aug. 23, 1989]

### § 3.372 Initial grant following inactivity of tuberculosis.

When service connection is granted initially on an original or reopened claim for pulmonary or nonpulmonary tuberculosis and there is satisfactory evidence that the condition was active previously but is now inactive (arrested), it will be presumed that the disease continued to be active for 1 year after the last date of established activity, provided there is no evidence to establish activity or inactivity in the intervening period. For a veteran entitled to receive compensation on August 19, 1968, the beginning date of graduated ratings will commence at the end of the 1-year period. For a veteran who was not receiving or entitled to receive compensation on August 19, 1968, ratings will be assigned in accordance with the Schedule for Rating Disabilities (part 4 of this chapter). This section is not applicable to running award cases.

[33 FR 16275, Nov. 6, 1968]

### § 3.373 [Reserved]

### § 3.374 Effect of diagnosis of active tuberculosis.

(a) *Service diagnosis.* Service department diagnosis of active pulmonary tuberculosis will be accepted unless a

board of medical examiners, Clinic Director or Chief, Outpatient Service certifies, after considering all the evidence, including the favoring or opposing tuberculosis and activity, that such diagnosis was incorrect. Doubtful cases may be referred to the Chief Medical Director in Central Office.

(b) *Department of Veterans Affairs diagnosis.* Diagnosis of active pulmonary tuberculosis by the medical authorities of the Department of Veterans Affairs as the result of examination, observation, or treatment will be accepted for rating purposes. Reference to the Clinic Director or Chief, Outpatient Service, will be in order in questionable cases and, if necessary, to the Chief Medical Director in Central Office.

(c) *Private physician's diagnosis.* Diagnosis of active pulmonary tuberculosis by private physicians on the basis of their examination, observation or treatment will not be accepted to show the disease was initially manifested after discharge from active service unless confirmed by acceptable clinical, X-ray or laboratory studies, or by findings of active tuberculosis based upon acceptable hospital observation or treatment.

[26 FR 1591, Feb. 24, 1961, as amended at 27 FR 6387, July 6, 1962; 33 FR 16275, Nov. 6, 1968; 43 FR 45348, Oct. 2, 1978]

### § 3.375 Determination of inactivity (complete arrest) in tuberculosis.

(a) *Pulmonary tuberculosis.* A veteran shown to have had pulmonary tuberculosis will be held to have reached a condition of "complete arrest" when a diagnosis of inactive is made.

(b) *Nonpulmonary disease.* Determination of complete arrest of nonpulmonary tuberculosis requires absence of evidence of activity for 6 months. If there are two or more foci of such tuberculosis, one of which is active, the condition will not be considered to be inactive until the tuberculous process has reached arrest in its entirety.

(c) *Arrest following surgery.* Where there has been surgical excision of the lesion or organ, the date of complete arrest will be the date of discharge from the hospital, or 6 months from the date of excision, whichever is later.

[33 FR 16275, Nov. 6, 1968, as amended at 43 FR 45348, Oct. 2, 1978]